



Thermodynamic Evaluation of the Production of Chiral Amines from Long-Chain Aliphatic Alcohols

Abu, Rohana; Lima Ramos, Joana; Woodley, John

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Abu, R., Lima Ramos, J., & Woodley, J. (2014). *Thermodynamic Evaluation of the Production of Chiral Amines from Long-Chain Aliphatic Alcohols*. Poster session presented at 3rd Multistep Enzyme Catalyzed Processes Congress, Madrid, Spain.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Thermodynamic Evaluation of the Production of Chiral Amines from Long-Chain Aliphatic Alcohols



Rohana Abu, Joana Lima-Ramos and John M. Woodley

Department of Chemical and Biochemical Engineering, Technical University of Denmark, Søtofts Plads Building 229, Kgs. Lyngby, Denmark. roha@kt.dtu.dk

Introduction


ω -Transaminase (EC.2.6.1.X) has been receiving considerable attention in chiral amine production due to its excellent stereoselectivity and its ability to operate under mild conditions. Potentially of even greater interest is to use alcohols as the starting material rather than ketones to produce the corresponding amines using a combination of an alcohol dehydrogenase (EC.1.1.1.2) in combination with the ω -transaminase [1, 2]. However of both reactions are reversible and often limited by thermodynamics where the equilibrium positions are in the favour of the reactants rather than the products.

Thermodynamics

Thermodynamic equilibrium of a system is very crucial it in large part responsible for determining the yield of the reaction. However, very little attention has been addressed on this issue in the scientific literature compared to the kinetic study of enzyme-catalysed reactions [3,4].

- For unfavourable reaction equilibrium, the reaction can be made possible and improved by shifting the equilibrium position towards the product with reaction process engineering and engineering strategies.
- For favourable reaction the study on kinetics will be useful and significant for control strategies.


Process Challenges



Solutions:

1. Substrate feeding strategy
2. Co-factor regeneration or enzyme cascade
3. Excess of amine donor
4. *In-situ* co-product removal or enzyme cascade
5. *In-situ* product removal
6. Biocatalyst format & amount
7. Reactor configuration

Example Flowsheet



Thermodynamic estimation

- Experimental methodology to determine K_{eq} [3]

$$Q = \frac{[C][D]}{[A][B]} \xrightarrow{t \rightarrow \infty} K$$

[A],[B],[C],[D] corresponds to reactant concentrations

$$\frac{Q_t}{Q_0} \rightarrow 1, \text{ when } Q \sim K$$

- Predictive tools to calculate of ΔG value using group contribution method [5, 6]

Process Perspective

There is a need to predict the thermodynamic equilibrium value of the system in order to guide decision making and to set a realistic target for scale-up.

The engineering tools used to assist process development

- Thermodynamic modelling
- Kinetic modeling

References

- [1] Sattler, J. H., Fuchs, M., Tauber, K., Mutti, F. G., Faber, K., Pfeffer, J., Haas, T. and Kroutil, W. (2012). *Angewandte Chemie*, 51(36): 9156–9.
- [2] Tauber, K., Fuchs, M., Sattler, J. H., Pitzer, J., Pressnitz, D., Koszelewski, D., Faber, K., Pfeffer, J., Haas, T. and Kroutil, W. (2013). *Chemistry*, 19(12): 4030–5.
- [3] Tufvesson, P., Jensen, J. S., Kroutil, W. and Woodley, J. M. (2012). *Biotechnology and Bioengineering*, 109(8): 2159–62.
- [4] Tufvesson, P., Lima-ramos, J., Jensen, J. S., Al-haque, N., Neto, W. and Woodley, J. M. (2011). *Biotechnology and Bioengineering*, 108(7): 1479–1493.
- [5] Jankowski, M. D., Henry, C. S., Broadbelt, L. J. and Hatzimanikatis, V. (2008). *Biophysical Journal*, 95(3), 1487–99.
- [6] Marrero, J. and Gani, R. (2001). *Fluid Phase Equilibria*, 183–184, 183–208.